

# Getting Back to TOLAC

---

Benjamin Dorton, M.D., FACOG

New Orleans, LA

# No Disclosures

---

# Objectives

---

- Review the history of Cesarean section and Trial of Labor after Cesarean Section (TOLAC)/Vaginal birth after Cesarean Section (VBAC)
- Review current and historical TOLAC and VBAC trends
- Review risks and benefits of TOLAC and VBAC
- Examine appropriate tools to employ in TOLAC counseling to offer direction for patient selection
- Discuss TOLAC methods and related clinical guidance

# History of Cesarean Section & TOLAC

---

# CESAREAN SECTION— A BRIEF HISTORY



Plate XIII from Sculietus' *Armenamentarium chirurgicum bipartitum*, 1666

A Brochure to Accompany an Exhibition  
on the History of Cesarean Section at the  
**NATIONAL LIBRARY OF MEDICINE**  
30 April 1993 - 31 August 1993

By Jane Eliot Sewell, Ph.D.  
for The American College of Obstetricians and Gynecologists




From Suetonius' *Lives of the Twelve Caesars*, 1506 woodcut.

# History of TOLAC

- 1916: Dr. Cragin declared: "once a cesarean section, always a cesarean section."
- 
- 1923: First reported Vaginal Birth after Cesarean Section was reported in medical literature.
  - 1945: Low transverse cesarean section became standard of care.
  - 1950: Dr. Eastman described a 30-percent post-cesarean vaginal delivery rate (2% uterine rupture & 10% maternal mortality rate) at Johns Hopkins Hospital.
  - 1960's: Dr. Pauerstein reviewed observational studies that suggested TOLAC was a reasonable option.

# History of TOLAC

- Electronic fetal monitoring
- Decline of vaginal breech delivery
- Operative vaginal delivery training
- Shorter labor duration expectations
- Repeat cesarean sections
- Safety of cesarean section
- Medical-legal concerns
- Convenience




# History of TOLAC

---

- 1980's: National Institutes of Health (NIH) Consensus Development Conference on Vaginal Birth After Cesarean (1981) was convened.

# History of TOLAC



Source: F. Gary Cunningham, Kenneth J. Landon, Jill S. Dache, Barbara L. Hellman, Catherine Y. Spring, Brian M. Casey. *Williams Obstetrics*, 10th Edition. Copyright © McGraw Hill. All rights reserved.

# History of TOLAC



The NEW ENGLAND  
JOURNAL of MEDICINE

CURRENT ISSUE ▾ SPECIALTIES ▾ TC

SOUNDING BOARD



## The Risks of Lowering the Cesarean-Delivery Rate

Authors: Benjamin P. Sachs, M.B., B.S., D.P.H., Cindy Kobelin, M.D., Mary Ames Castro, M.D., and Fredric Frigoletto  
M.D. [Author Info & Affiliations](#)

Published January 7, 1999 | N Engl J Med 1999;340:54-57 | DOI: 10.1056/NEJM199901073400112

[VOL. 340 NO. 1](#) | [Copyright © 1999](#)

## ONCE A CESAREAN, ALWAYS A CONTROVERSY

*Bruce L. Flamm, MD*

For most of this century, “once a cesarean, always a cesarean” was the rule in the United States. In the 1980s, vaginal birth after cesarean grew in popularity and the pendulum began to swing away from routine repeat cesarean delivery. Recently, the wisdom of this transition has been questioned. As the 20th century comes to a close, the treatment of the patient with a prior cesarean delivery remains controversial. (Obstet Gynecol 1997;90:312–5. © 1997 by The American College of Obstetricians and Gynecologists.)

# History of TOLAC



The NEW ENGLAND  
JOURNAL of MEDICINE

CURRENT ISSUE ▾ SPECIALTIES ▾ TOPICS

ORIGINAL ARTICLE



## Comparison of a Trial of Labor with an Elective Second Cesarean Section

**Authors:** Michael J. McMahon, M.D., M.P.H., Edwin R. Luther, M.D., Watson A. Bowes, Jr., M.D., and Andrew F. Olshan, Ph.D. [Author Info & Affiliations](#)


Published September 5, 1996 | N Engl J Med 1996;335:689-695 | DOI: 10.1056/NEJM199609053351001

[VOL. 335 NO. 10](#) | [Copyright © 1996](#)

### CONCLUSIONS


Among pregnant women who have had a cesarean section, major maternal complications are almost twice as likely among those whose deliveries are managed with a trial of labor as among those who undergo an elective second cesarean section.

ACOG PB No. 5, 1999: Vaginal birth after previous cesarean delivery



**Fig. 1.** Cesarean delivery rates from 1970 to 2009. (Data from <http://www.cdc.gov/nchs/nvss.htm>, courtesy of Caroline Signore, MD, MPH.)

*Scott. VBAC: A Common-Sense Approach. Obstet Gynecol 2011.*



**Fig. 2.** Rates of total cesarean deliveries (1998–2009), primary cesarean deliveries (1998–2007), and vaginal births after cesarean (VBAC) (1998 to 2007). (Data from <http://www.cdc.gov/nchs/nvss.htm>, courtesy of Caroline Signore, MD, MPH.)

*Scott. VBAC: A Common-Sense Approach. Obstet Gynecol 2011.*

# History of TOLAC

Review

> NIH Consens State Sci Statements. 2010 Mar 10;27(3):1-42.

## NIH consensus development conference draft statement on vaginal birth after cesarean: new insights




F Gary Cunningham <sup>1</sup>, Shrikant I Bangdiwala, Sarah S Brown, Thomas Michael Dean, Marilyn Frederiksen, Carol J Rowland Hogue, Tekoa King, Emily Spencer Lukacz, Laurence B McCullough, Wanda Nicholson, Nancy Frances Petit, Jeffrey Lynn Probstfield, Adele C Viguera, Cynthia A Wong, Sheila Cohen Zimmet

# History of TOLAC

- Main 2010 panel conclusions:
- Trial of labor is a **reasonable option** for many pregnant individuals with **one prior low transverse uterine incision**.
- One of the panel's major goals is to support pregnant individuals with one prior transverse uterine incision to **make informed decisions** about trial of labor compared with elective repeat cesarean delivery.
- When trial of labor and elective repeat cesarean delivery are medically equivalent options, a **shared decision making process should be adopted** and, whenever possible, the patient's preference should be honored.
- Facilities should **improve access for TOLAC**.
- **Medical-legal considerations** add to, and in many instances exacerbate, these barriers to trial of labor.

# History of TOLAC



Source: F. Gary Cunningham, Kenneth J. Leveno, Juli S. Dwyer, Barbara L. Hoffman, Catherine Y. Spring, Brian M. Casey. Williams Obstetrics, 24th Edition. Copyright © McGraw Hill. All rights reserved.

# To TOLAC or Not to TOLAC?

---

# Who are not candidates for TOLAC?

---

- Prior classical or T uterine incision (risk of rupture 4-9%)
- Prior uterine rupture (risk of rupture ~10%)
- Prior myomectomy with entry into endometrium or full thickness myometrial disruption
- Placenta previa or accreta
- Vasa previa
- Active genital herpes
- Fetal malpresentation
- Cord prolapse

# Risk Stratification for TOLAC candidates

---

- Increased success rate
  - Nonrecurring indication for first cesarean delivery
  - Spontaneous labor
  - Younger maternal age
  - Low BMI
  - Gestational age  $\leq$  40 weeks
  - Normal birth weight
  - Prior vaginal delivery
- Decreased success rate
  - Recurring indication for first cesarean delivery
  - Induction of labor
  - Older maternal age
  - Higher BMI
  - Gestational age  $>$  40 weeks
  - Higher birth weight/Macrosomia
  - Short interdelivery interval ( $<$ 19 months)
  - Preeclampsia

# To TOLAC or Not to TOLAC?

---

- MFMU VBAC calculator
- No RCT exist for maternal or neonatal outcomes for TOLAC vs. Repeat C/S

## VAGINAL BIRTH AFTER CESAREAN

Early Pregnancy

Delivery Admission

Maternal age (range 15-50 years):

Height Unit:

- inches  
 centimeters

Height (range 46-75 in):

Weight Unit:

- pounds  
 kilograms

Pre-pregnancy weight (range 74-454 lbs):

Body mass index: kg/m<sup>2</sup>

Obstetric History:

Previous VBAC

Arrest disorder indication for prior cesarean?


No

Treated chronic hypertension?

No

Calculate

# Benefits of successful TOLAC



# Risks of TOLAC

**Table 1.** Composite Maternal Risks From Elective Repeat Cesarean Delivery and Trial of Labor After Previous Cesarean Delivery in Term Patients

Maternal Risks	ERCD (%) [One CD]	TOLAC (%)
Infectious morbidity	3.2	4.6
Surgical injury	0.30–0.60	0.37–1.3
Blood transfusion	0.46	0.66
Hysterectomy	0.16	0.14
Uterine rupture	0.02	0.71
Maternal death	0.0096	0.0019

Abbreviations: CD, cesarean delivery; ERCD, elective repeat cesarean delivery; TOLAC, trial of labor after cesarean delivery.

**Surgical Injury:** Defined differently and variably reported on in trials. Rate of surgical injury may be increased with TOLAC but definitive studies are lacking.

**Infectious Morbidity:** Defined as fever, infection, endometritis, and chorioamnionitis

Data from Guise JM, Eden K, Emeis C, Denman MA, Marshall N, Fu R, et al. Vaginal birth after cesarean: new insights. (Archived) Evidence Report/Technology Assessment No.191. AHRQ Publication No. 10-E003. Rockville (MD): Agency for Healthcare Research and Quality; 2010.

# Risks of TOLAC


**Table 2.** Composite Neonatal Morbidity From Elective Repeat Cesarean Delivery and Trial of Labor After Previous Cesarean Delivery in Term Infants

Neonatal Risks	ERCD (%)	TOLAC (%)
Antepartum stillbirth	0.21	0.10
Intrapartum stillbirth	0–0.004	0.01–0.04
HIE	0–0.32	0–0.89
Perinatal mortality	0.05	0.13
Neonatal mortality	0.06	0.11
NICU admission	1.5–17.6	0.8–26.2
Respiratory morbidity	2.5	5.4
Transient tachypnea	4.2	3.6

Abbreviations: ERCD, elective repeat cesarean delivery; HIE, hypoxic ischemic encephalopathy; NICU, neonatal intensive care unit; TOLAC, trial of labor after cesarean delivery.

# To TOLAC or Not to TOLAC?

---



# To TOLAC or Not to TOLAC?

---



Cesarean  
deliveries



PAS



# To TOLAC or Not to TOLAC?

---


60-70%  
likelihood  
of VBAC  
success



Similar  
maternal  
morbidity  
as ERCS

# To TOLAC or Not to TOLAC?

---



# To TOLAC or Not to TOLAC?

Patient's with more than one cesarean section

## Risk of Uterine Rupture With a Trial of Labor in Women With Multiple and Single Prior Cesarean Delivery

*Mark B. Landon, MD, Catherine Y. Spong, MD, Elizabeth Thom, PhD, John C. Hauth, MD, Steven L. Bloom, MD, Michael W. Varner, MD, Atef H. Moawad, MD, Steve N. Caritis, MD, Margaret Harper, MD, MS, Ronald J. Wapner, MD, Yoram Sorokin, MD, Menachem Miodovnik, MD, Marshall Carpenter, MD, Alan M. Peaceman, MD, Mary J. O'Sullivan, MD, Baha M. Sibai, MD, Oded Langer, MD, John M. Thorp, MD, Susan M. Ramin, MD, Brian M. Mercer, MD, and Steven G. Gabbe, MD, for the National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network\**

Obstetrics and Gynecology, Vol 108, No. 1, July 2006

# To TOLAC or Not to TOLAC?

Patient's with more than one cesarean section

Table 2. Maternal and Perinatal Outcomes

Outcome	Multiple (n = 975)	Single (n = 16,915)	OR (95% CI)	P
Uterine rupture	9 (0.9)	115 (0.7)	1.36 (0.69–2.69)	.37
Endometritis	30 (3.1)	485 (2.9)	1.08 (0.74–1.56)	.70
Hysterectomy	6 (0.6)	35 (0.2)	2.99 (1.25–7.12)	.023
Transfusion	31 (3.2)	273 (1.6)	2.00 (1.37–2.92)	< .001
Thromboembolic disease*	1 (0.1)	6 (0.04)	2.90 (0.35–24.09)	.32
Operative injury†	4 (0.4)	60 (0.4)	1.16 (0.42–3.19)	.78
Maternal death	0 (0.0)	3 (0.02)	–	1.00
Maternal composite‡	71 (7.3)	829 (4.9)	1.53 (1.19–1.96)	.001
Term NICU admission§	75 (11.2)	1321 (9.0)	1.28 (1.00–1.63)	.05
Term intrapartum stillbirth§	0 (0.0)	2 (0.01)	–	1.00
Term neonatal death§	1 (0.15)	12 (0.08)	1.83 (0.24–14.08)	.44
Term HIE§	0 (0.0)	12 (0.1)	–	3b.00

# To TOLAC or Not to TOLAC?


Patient's with more than one cesarean section

Outcome	TOLAC following 2 previous CS	ELCS following 2 CS	TOLAC following 1 CS	P- value
	Mean ( $\pm$ SD) or N (%)	Mean ( $\pm$ SD) or N (%)	Mean ( $\pm$ SD) or N (%)	
	N = 146	N = 206	N = 99	
Uterine rupture	1 (0.7 %)	N/A	0 (0 %)	1
Endometritis/ sepsis*	15 (10.3 %)	1 (0.5 %)	3 (3 %)	<0.01
Length of hospital stay (days)*	2.4 ( $\pm$ 1.8)	1.8 ( $\pm$ 0.8)	1.8 ( $\pm$ 1.7)	<0.01
Apgar at 5 min (score < 7)	8 (5.5 %)	1 (0.49 %)	1 (1 %)	<0.01
NIU admission	20 (13.7 %)	11 (5.3 %)	4 (4.0 %)	<0.01

# To TOLAC or Not to TOLAC?

Patient's with twin gestation

Studies have demonstrated similar outcomes to singleton gestations without increased risk





American Journal of Obstetrics and Gynecology

Volume 193, Issue 3, Supplement, September 2005, Pages 1050-1055




Transactions of the Twenty-Fifth Annual Meeting of the Society for Maternal-Fetal Medicine

## Vaginal birth after cesarean (VBAC) attempt in twin pregnancies: Is it safe?

Alison Cahill MD <sup>a</sup>  , David M. Stamilio MD, MSCE <sup>a</sup>, Emmanuelle Paré MD, FRCSC <sup>a b</sup>, Jeffrey P. Peipert MD, MPH <sup>e</sup>, Erika J. Stevens MS <sup>a b</sup>, Deborah B. Nelson PhD <sup>b c</sup>, George A. Macones MD, MSCE <sup>a b c d</sup>

# Let's TOLAC



# Let's TOLAC

---

- -Spontaneous labor is associated with an increased likelihood of achieving a VBAC.
- Induction of labor at 39-41 weeks gestation is associated with a lower odds of cesarean section when compared to expectant management

# Let's TOLAC

## Fetal monitoring

- External continuous monitoring

## Cervical ripening

- Oxytocin and mechanical but avoid misoprostol

## External cephalic version


- Not contraindicated
- Similar success rates

## Epidural

- Shared decision making

## Location

- Level 1 hospital or higher
- Ability to perform emergent deliveries




# Uterine rupture


## Uterine Rupture Trends in Patients Pursuing Trial of Labor After Cesarean in the United States from 2010 to 2022

Lillian H. Goodman, MD, MPH, Amanda A. Allshouse, MS, Torri D. Metz, MD, MS, and Ann M. Bruno, MD, MS  
Obstetrics and Gynecology. Vol. 145, No. 5, May 2025.

- Overall uterine rupture rate was 0.28%.
- Annual increase of 6.2% per year.




A





B

# Vaginal birth after Cesarean deliveries: United States, 2019-2024

Rate per 100 births to women with a previous Cesarean



© 2026 March of Dimes. All rights reserved.



Source: A. Rice, Contraception, 70(1), 2002, pp. 1-10. © 2002, Elsevier. Reprinted with permission. © 2002, Elsevier. Reprinted with permission. © 2002, Elsevier. Reprinted with permission.

# AI for TOLAC Prediction?

---

- Current standard remains Grobman VBAC calculator
- Some emerging AI models include Random Forest, XGBoost, and Neural Networks
- Early studies suggest XGBoost may offer superior discrimination compared to traditional logistic regression
- However, widespread adoption awaits prospective validation, interpretability, and integration into clinical workflows

# Summary

---

- Patient's with one or two prior low transverse cesarean delivery should be counseled and offered TOLAC.
- The overall rate of maternal and neonatal morbidity is low for patient's attempting TOLAC.
- Delivery should occur at a level one facility or higher with the ability to perform emergent deliveries.
- Acceptable induction agents during TOLAC include mechanical and oxytocin but Misoprostol is contraindicated.
- Evidence is based on retrospective and observational studies but no RCT comparing maternal and neonatal outcomes exists.

THANK YOU

# References

1. Cesarean section-A brief history. A Brochure to accompany an exhibition on the history of cesarean section at the National library of medicine April 30, 1993- August 31, 1993. Jane Eliot Sewell for ACOG.
2. Alavi-Tabari N, Black A. Trial of Labor After Previous Cesarean Section. In: Posner GD, Black AY, Jones G, El-Chaâr D. eds. *Oxorn-Foote Human Labor & Birth, 7e*. Chapter 23. McGraw-Hill Education; 2022.
3. Prior Cesarean Delivery." *Williams Obstetrics, 23e* Eds. F. Gary Cunningham, et al. McGraw Hill, 2010.
4. Prior Cesarean Delivery." *Williams Obstetrics, 26e* Eds. F. Gary Cunningham, et al. McGraw Hill, 2022.
5. The cesarean birth epidemic: Trends, causes, and solutions. Porreco, Richard P. et al. *American Journal of Obstetrics & Gynecology*, Volume 175, Issue 2, 369 – 374.
6. Scott JR. Vaginal birth after cesarean delivery: a common-sense approach. *Obstet Gynecol.* 2011 Aug;118(2 Pt 1):342-350.
7. Sachs BP, Kobelin C, Castro MA, Frigoletto F. The risks of lowering the cesarean-delivery rate. *N Engl J Med.* 1999 Jan 7;340(1):54-7
8. ACOG practice bulletin. Vaginal birth after previous cesarean delivery. Number 5, July 1999.
9. Flamm BL. Once a cesarean, always a controversy. *Obstet Gynecol.* 1997 Aug;90(2):312-5.
10. McMahon MJ, Luther ER, Bowes WA Jr, Olshan AF. Comparison of a trial of labor with an elective second cesarean section. *N Engl J Med.* 1996 Sep 5;335(10):689-95
11. Cunningham FG, Bangdiwala SI, Brown SS, Dean TM, Frederiksen M, Rowland Hogue CJ, King T, Spencer Lukacz E, McCullough LB, Nicholson W, Petit NF, Probstfield JL, Viguera AC, Wong CA, Zimmet SC. NIH consensus development conference draft statement on vaginal birth after cesarean: new insights. *NIH Consens State Sci Statements.* 2010 Mar 10;27(3):1-42.
12. Landon MB, Spong CY, Thom E, Hauth JC, Bloom SL, Varner MW, Moawad AH, Caritis SN, Harper M, Wapner RJ, Sorokin Y, Miodovnik M, Carpenter M, Peaceman AM, O'sullivan MJ, Sibai BM, Langer O, Thorp JM, Ramin SM, Mercer BM, Gabbe SG; National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Risk of uterine rupture with a trial of labor in women with multiple and single prior cesarean delivery. *Obstet Gynecol.* 2006 Jul;108(1):12-20.
13. J.C. McMullan et al. Trial of labour following two previous caesarean sections – A UK cohort study. *European Journal of Obstetrics & Gynecology and Reproductive Biology* 298 (2024) 182–186
14. Sugai S, Yamawaki K, Haino K, Yoshihara K, Nishijima K. Incidence of Recurrent Uterine Rupture: A Systematic Review and Meta-analysis. *Obstet Gynecol.* 2023 Dec 1;142(6):1365-1372.
15. Varner MW, Thom E, Spong CY, Landon MB, Leveno KJ, Rouse DJ, Moawad AH, Simhan HN, Harper M, Wapner RJ, Sorokin Y, Miodovnik M, Carpenter M, Peaceman A, O'sullivan MJ, Sibai BM, Langer O, Thorp JM, Ramin SM, Mercer BM; National Institute of Child Health and Human Development (NICHD) Maternal-Fetal Medicine Units Network (MFMU). Trial of labor after one previous cesarean delivery for multifetal gestation. *Obstet Gynecol.* 2007 Oct;110(4):814-9.
16. Cahill A, Stamilio DM, Paré E, Peipert JP, Stevens EJ, Nelson DB, Macones GA. Vaginal birth after cesarean (VBAC) attempt in twin pregnancies: is it safe? *Am J Obstet Gynecol.* 2005 Sep;193(3 Pt 2):1050-5.
17. Goodman LH, Allshouse AA, Metz TD, Bruno AM. Uterine Rupture Trends in Patients Pursuing Trial of Labor After Cesarean in the United States from 2010 to 2022. *Obstet Gynecol.* 2025 May 1;145(5):511-513. doi: 10.1097/AOG.0000000000005882. Epub 2025 Mar 13